

Daily Announcements & Reminders:**Goals for Today:**

Section 16.7/16.8

- Apply Stokes' Theorem to flux integral problems.
- Use Stokes' Theorem to simplify flux integrals
- Introduce and apply the Divergence Theorem to flux integral problems

Theorem 136 (Stokes' Theorem). *Let S be a smooth oriented surface and C be its compatibly oriented boundary. Let \mathbf{F} be a vector field with continuous partial derivatives. Then*

$$\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\sigma = \int_C \mathbf{F} \cdot \mathbf{T} \, ds.$$

Example 137 (DD). Let $\mathbf{F} = \langle -y, x + (z-1)x^{x \sin(x)}, x^2 + y^2 \rangle$. Find $\iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\sigma$ over the surface S which is the part of the sphere $x^2 + y^2 + z^2 = 2$ above $z = 1$, oriented away from the origin.

Question: What can we say if two different surfaces S_1 and S_2 have the same oriented boundary C ?

Example 138. Suppose $\text{curl } \mathbf{F} = \langle y^{y^y} \sin(z^2), (y-1)e^{x^{x^x}} + 2, -ze^{x^{x^x}} \rangle$. Compute the net flux of the curl of \mathbf{F} over the surface pictured below, which is oriented outward and whose boundary curve is a unit circle centered on the y -axis in the plane $y = 1$.

Theorem 139 (Divergence Theorem). *Let S be a closed surface oriented outward, D be the volume inside S , and \mathbf{F} be a vector field with continuous partial derivatives. Then*

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, d\sigma = \iiint_D \nabla \cdot \mathbf{F} \, dV.$$

Example 140. Let $\mathbf{F} = \langle y^{1234}e^{\sin(yz)}, y - x^{z^x}, x^2 - z \rangle$ and S be the surface consisting of the portion of cylinder of radius 1 centered on the z -axis between $z = 0$ and $z = 3$, together with top and bottom disks, oriented outward. Find the flux of \mathbf{F} through S .

