

**MATH 2551-D MIDTERM 1**  
**VERSION A**  
**SPRING 2023**  
**COVERS SECTIONS 12.1-12.6, 13.1-13.4**

**Full name:** \_\_\_\_\_

**GT ID:** \_\_\_\_\_

**Honor code statement:** I will abide strictly by the Georgia Tech honor code at all times. I will not use a calculator. I will not reference any website, application, or other CAS-enabled service. I will not consult with my notes or anyone during this exam. I will not provide aid to anyone else during this exam.

(     ) I attest to my integrity.

**Read all instructions carefully** before beginning.

- Print your name and GT ID neatly above.
- You have 75 minutes to take the exam.
- You may not use aids of any kind.
- Please show your work.
- Good luck! Write yourself a message of encouragement on the front page!

Question	Points
1	2
2	2
3	2
4	4
5	10
6	10
7	10
8	10
Total:	50



For problems 1-3 choose whether each statement is true or false. If the statement is *always* true, pick true. If the statement is *ever* false, pick false. Be sure to neatly fill in the bubble corresponding to your answer choice.

1. (2 points) The curvature of a line is 0 at every point on the line.

**TRUE**

**FALSE**

2. (2 points) If  $\mathbf{u}$  and  $\mathbf{v}$  are two vectors in  $\mathbb{R}^3$ , then  $\mathbf{u} \cdot \mathbf{v}$  is a vector orthogonal to both  $\mathbf{u}$  and  $\mathbf{v}$ .

**TRUE**

**FALSE**

3. (2 points) There is exactly one possible parameterization for any line in space.

**TRUE**

**FALSE**

4. (4 points) Which of the following is true of the quadric surface  $z = x^2 + 2y^2$ ?

**A)** It is a plane, because all of its cross-sections in the  $x = k$ ,  $y = k$ , and  $z = k$  planes are straight lines.

**B)** It is a sphere, because all of its cross-sections in the  $x = k$ ,  $y = k$ , and  $z = k$  planes are circles.

**C)** It is an elliptical paraboloid, because its cross-sections are ellipses in the  $z = k$  planes and parabolas in the  $x = k$  and  $y = k$  planes.

**D)** It is a hyperbolic paraboloid, because its cross-sections are hyperbolas in the  $z = k$  planes and parabolas in the  $x = k$  and  $y = k$  planes.

**E)** It is a cone, because its cross-sections are circles in the  $z = k$  planes and straight lines in the  $x = k$  and  $y = k$  planes.

5. Let  $C$  be the curve which is the graph of the vector-valued function  $\mathbf{r}(t) = \left\langle \frac{t^2}{2}, \frac{4}{3}t^{3/2}, 2t \right\rangle$ ,  $0 \leq t < \infty$ .

(a) (2 points) Compute  $\mathbf{r}(0)$  and  $\mathbf{r}(4)$ .

(b) (6 points) Compute the arc length function  $s(t)$  for this parameterization, taking  $t_0 = 0$  as your reference point.

(c) (2 points) Find the distance along  $C$  between the points  $(0, 0, 0)$  and  $(8, \frac{32}{3}, 8)$ .

6. Suppose that a pegasus (a flying horse) is flying through the sky. Two hours after it left its stable, its velocity vector is  $\mathbf{v} = 3\mathbf{i} - 2\mathbf{j} + \mathbf{k}$  miles per hour and its position is  $2\mathbf{i} - 3\mathbf{j} + 4\mathbf{k}$  miles. The stable is not located at the origin.
- (a) (3 points) Find an equation of the tangent line to the flight path of the pegasus two hours after it left its stable.
- (b) (5 points) One hour after it left its stable, the position of the pegasus was  $-\mathbf{i} - \mathbf{j} + 5\mathbf{k}$  miles. Use this fact and your answer to part (a) to show that the pegasus cannot be flying in a straight line. Be sure to clearly explain your answer.
- (c) (2 points) If the pegasus maintains its flight speed at the two hour mark of  $\sqrt{14}$  miles per hour for the rest of its flight, is it possible for it to return to its stable? Be sure to clearly explain your answer.

7. Let  $P_1$  be the plane  $2x + 2y - z = 1$  and  $P_2$  be the plane  $2x - y + 2z = 4$ .

(a) (2 points) Show that  $P_1$  and  $P_2$  are intersecting planes by showing that the point  $(-1, 4, 5)$  lies on both planes.

(b) (2 points) Find normal vectors to both planes.

(c) (6 points) Find a parameterization of the line of intersection of the two planes.

8. In this problem, you will work with normal vectors and curvature.

(a) (4 points) Let  $\mathbf{r}(t) = f(t)\mathbf{i} + g(t)\mathbf{j}$  be any parameterized curve in the plane. Show that the vector  $\mathbf{n}(t) = -g'(t)\mathbf{i} + f'(t)\mathbf{j}$  is orthogonal to  $\mathbf{r}'(t)$  for every value of  $t$ .

(b) (4 points) Use your result from part (a) to produce a **unit** normal vector to the curve  $\mathbf{r}(t) = \langle 2\cos(t), 2\sin(t) \rangle, 0 \leq t < 2\pi$ .

(c) (2 points) Is the vector that you produced in part (b) the principal unit normal vector? Explain your answer; a picture might be useful.  
*Hint: Consider whether your vector is pointing in the direction of curvature of this curve.*

**SCRATCH PAPER - PAGE LEFT BLANK**



## FORMULA SHEET

- $\langle u_1, u_2, u_3 \rangle \cdot \langle v_1, v_2, v_3 \rangle = u_1v_1 + u_2v_2 + u_3v_3$

- $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}||\mathbf{v}| \cos(\theta)$

- $\langle u_1, u_2, u_3 \rangle \times \langle v_1, v_2, v_3 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$

- $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}||\mathbf{v}| |\sin(\theta)|$

- $L = \int_a^b |\mathbf{r}'(t)| dt$

- $s(t) = \int_{t_0}^t |\mathbf{r}'(T)| dT$

- $\mathbf{T} = \frac{\mathbf{v}}{|\mathbf{v}|} = \frac{d\mathbf{r}}{ds}$

- $\kappa = \left| \frac{d\mathbf{T}}{ds} \right| = \frac{1}{|\mathbf{v}|} \left| \frac{d\mathbf{T}}{dt} \right| = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3}$

- $\mathbf{N} = \frac{1}{\kappa} \frac{d\mathbf{T}}{ds} = \frac{d\mathbf{T}/dt}{|d\mathbf{T}/dt|}$

**SCRATCH PAPER - PAGE LEFT BLANK**