

MATH 2551-K FINAL EXAM
PART 1
VERSION C
FALL 2023
COVERS SECTIONS 12.1-12.6, 13.1-13.4, 14.1

Full name: _____

GT ID: _____

Honor code statement: I will abide strictly by the Georgia Tech honor code at all times. I will not use a calculator. I will not reference any website, application, or other CAS-enabled service. I will not consult with my notes or anyone during this exam. I will not provide aid to anyone else during this exam.

() I attest to my integrity and will not discuss this exam with anyone until **Friday December 15**.

Read all instructions carefully before beginning.

- Print your name and GT ID neatly above.
- You may use any portion of our 2 hours and 50 minute final exam time to work on this portion of the exam.
- When you complete this portion of the exam, you may turn it in and work on another portion of the exam.
- You may not use aids of any kind.
- Please show your work.
- Good luck! Write yourself a message of encouragement on the front page!

Question	Points
1	2
2	2
3	3
4	3
5	10
6	10
7	10
Total:	40

For problems 1-3 choose whether each statement is true or false. If the statement is *always* true, pick true. If the statement is *ever* false, pick false. Be sure to neatly fill in the bubble corresponding to your answer choice.

1. (2 points) If \mathbf{u} and \mathbf{v} are unit vectors in \mathbb{R}^3 , then $|\mathbf{u} \times \mathbf{v}| = 1$.

TRUE

FALSE

2. (2 points) All surfaces in \mathbb{R}^3 are quadric surfaces.

TRUE

FALSE

3. (3 points) Level surfaces of the function $f(x, y, z) = \sqrt{x^2 + z^2}$ are:

- Circles centered at the origin
- Spheres centered at the origin
- Upper hemispheres centered at the origin
- Cylinders centered around the y -axis
- None of the above

4. (3 points) Which of the following planes is parallel to the plane $y = -2 - 4x + 3z$?

- $x + y + 3z = 1$
- $8x + 2y - 6z = -5$
- $4x + y + 3z = -2$
- $42x + 3z = -2$
- None of the above

5. (10 points) Find the length of the portion of the helix $\mathbf{r}(t) = \langle 3 \sin(t), 4t, 3 \cos(t) \rangle$ between $(0, 0, 3)$ and $(3, 2\pi, 0)$.

6. (a) (3 points) Find an equation of the plane perpendicular to $\mathbf{n} = \mathbf{j} + \mathbf{k}$ that passes through the point $P = (1, 2, 3)$.

(b) (3 points) Find an equation of the plane perpendicular to the line $\ell(t) = \langle 3, t-1, t-17 \rangle$ that passes through the point $Q = (2, 3, 4)$.

(c) (4 points) Use your work in parts (a) and (b) to explain why there is no plane with normal vector parallel to $\langle 0, 0, 1 \rangle$ that contains both the points P and Q .

7. Consider the curve parameterized by $\mathbf{r}(t) = \langle \sqrt{23}e^t, e^t \cos(t), e^t \sin(t) \rangle$ for $t \in \mathbb{R}$.

(a) (5 points) Compute the unit tangent vector $\mathbf{T}(t)$.

(b) (5 points) Compute the curvature $\kappa(t)$.

SCRATCH PAPER - PAGE LEFT BLANK

FORMULA SHEET

- $\langle u_1, u_2, u_3 \rangle \cdot \langle v_1, v_2, v_3 \rangle = u_1v_1 + u_2v_2 + u_3v_3$

- $\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}||\mathbf{v}|\cos(\theta)$

- $\langle u_1, u_2, u_3 \rangle \times \langle v_1, v_2, v_3 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$

- $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}||\mathbf{v}|\sin(\theta)$

- $L = \int_a^b |\mathbf{r}'(t)| dt$

- $s(t) = \int_{t_0}^t |\mathbf{r}'(T)| dT$

- $\mathbf{T} = \frac{\mathbf{v}}{|\mathbf{v}|} = \frac{d\mathbf{r}}{ds}$

- $\kappa = \left| \frac{d\mathbf{T}}{ds} \right| = \frac{1}{|\mathbf{v}|} \left| \frac{d\mathbf{T}}{dt} \right| = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3}$

- $\mathbf{N} = \frac{1}{\kappa} \frac{d\mathbf{T}}{ds} = \frac{d\mathbf{T}/dt}{|d\mathbf{T}/dt|}$

SCRATCH PAPER - PAGE LEFT BLANK

MATH 2551-K FINAL EXAM
PART 2
VERSION C
FALL 2023
COVERS SECTIONS 14.2-14.8, 15.1-15.4

Full name: _____

GT ID: _____

Honor code statement: I will abide strictly by the Georgia Tech honor code at all times. I will not use a calculator. I will not reference any website, application, or other CAS-enabled service. I will not consult with my notes or anyone during this exam. I will not provide aid to anyone else during this exam.

() I attest to my integrity and will not discuss this exam with anyone until **Friday December 15**.

Read all instructions carefully before beginning.

- Print your name and GT ID neatly above.
- You may use any portion of our 2 hours and 50 minute final exam time to work on this portion of the exam.
- When you complete this portion of the exam, you may turn it in and work on another portion of the exam.
- You may not use aids of any kind.
- Please show your work.
- Good luck! Write yourself a message of encouragement on the front page!

Question	Points
1	2
2	2
3	3
4	3
5	10
6	10
7	10
Total:	40

For problems 1-3 choose whether each statement is true or false. If the statement is *always* true, pick true. If the statement is *ever* false, pick false. Be sure to neatly fill in the bubble corresponding to your answer choice.

1. (2 points) The total derivative of a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}^7$ at the point (a, b, c) is a 2×7 matrix.

TRUE

FALSE

2. (2 points) Any surface that is a graph of a function of two variables $z = f(x, y)$ can be thought of as a level surface of a function of 3 variables.

TRUE

FALSE

3. (3 points) Compute the rate of change of the function $f(x, y, z) = 3xy + z^2$ at the point $(3, 0, 3)$ in the direction of $\langle 2, 1, 2 \rangle$.

5

7

15

21

None of the above.

4. (3 points) Find the linearization of the function $f(x, y) = \sqrt{x^2 + y}$ at the point $(2, 5)$.

$L(x, y) = 3 + \frac{2}{3}(x - 2) + \frac{1}{6}(y - 5)$

$L(x, y) = \frac{2}{3}(x - 2) + \frac{1}{6}(y - 5)$

$L(x, y) = \frac{1}{6}(x - 2) + \frac{1}{6}(y - 5)$

$L(x, y) = 3 + \frac{x}{\sqrt{x^2 + y}}(x - 2) + \frac{1}{2\sqrt{x^2 + y}}(y - 5)$

None of the above.

5. (10 points) Find and classify the critical points of the function $f(x, y) = 3x^3 + 3xy + 3y^3$.

6. For each limit below, either compute its value or show that it does not exist.


(a) (2 points) $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + 3y^4 + 1}$

(b) (4 points) $\lim_{(x,y) \rightarrow (0,0)} x \frac{x^2 y^2}{x^2 + 3y^4}$

Hint: This limit exists. Try converting to polar coordinates and taking the limit as $r \rightarrow 0$

(c) (4 points) $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + 3y^4}$

7. (10 points) The region R bounded by $y = e^x - 1$, $y = 2$, and $y = x/2$ is shown to the right. Write an iterated integral or sum of iterated integrals for the double integral $\iint_R e^x dA$, using either order of integration, then compute your integral. Fully simplify your answer.



SCRATCH PAPER - PAGE LEFT BLANK

FORMULA SHEET

- Total Derivative: For $\mathbf{f}(x_1, \dots, x_n) = \langle f_1(x_1, \dots, x_n), \dots, f_m(x_1, \dots, x_n) \rangle$

$$D\mathbf{f} = \begin{bmatrix} (f_1)_{x_1} & (f_1)_{x_2} & \cdots & (f_1)_{x_n} \\ (f_2)_{x_1} & (f_2)_{x_2} & \cdots & (f_2)_{x_n} \\ \vdots & \ddots & \cdots & \vdots \\ (f_m)_{x_1} & (f_m)_{x_2} & \cdots & (f_m)_{x_n} \end{bmatrix}$$

- Linearization: Near \mathbf{a} , $L(\mathbf{x}) = f(\mathbf{a}) + Df(\mathbf{a})(\mathbf{x} - \mathbf{a})$
- Chain Rule: If $h = g(f(\mathbf{x}))$ then $Dh(\mathbf{x}) = Dg(f(\mathbf{x}))Df(\mathbf{x})$
- Implicit Differentiation: $\frac{\partial z}{\partial x} = \frac{-F_x}{F_z}$ and $\frac{\partial z}{\partial y} = \frac{-F_y}{F_z}$.
- Directional Derivative: If \mathbf{u} is a unit vector, $D_{\mathbf{u}}f(P) = Df(P)\mathbf{u} = \nabla f(P) \cdot \mathbf{u}$
- The tangent line to a level curve of $f(x, y)$ at (a, b) is $0 = \nabla f(a, b) \cdot \langle x - a, y - b \rangle$
- The tangent plane to a level surface of $f(x, y, z)$ at (a, b, c) is

$$0 = \nabla f(a, b, c) \cdot \langle x - a, y - b, z - c \rangle.$$

- Hessian Matrix: For $f(x, y)$, $Hf(x, y) = \begin{bmatrix} f_{xx} & f_{yx} \\ f_{xy} & f_{yy} \end{bmatrix}$
- Second Derivative Test: If (a, b) is a critical point of $f(x, y)$ then
 1. If $\det(Hf(a, b)) > 0$ and $f_{xx}(a, b) < 0$ then f has a local maximum at (a, b)
 2. If $\det(Hf(a, b)) > 0$ and $f_{xx}(a, b) > 0$ then f has a local minimum at (a, b)
 3. If $\det(Hf(a, b)) < 0$ then f has a saddle point at (a, b)
 4. If $\det(Hf(a, b)) = 0$ the test is inconclusive

- Area/volume: $\text{area}(R) = \iint_R dA$, $\text{volume}(D) = \iiint_D dV$

- Trig identities: $\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$, $\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$

- Average value: $f_{\text{avg}} = \frac{\iint_R f(x, y) dA}{\text{area of } R}$

- Polar coordinates: $x = r \cos(\theta)$, $y = r \sin(\theta)$, $dA = r dr d\theta$

SCRATCH PAPER - PAGE LEFT BLANK

MATH 2551-K FINAL EXAM
PART 3
VERSION C
FALL 2023
COVERS SECTIONS 15.1-15.8, 16.1-16.8

Full name: _____

GT ID: _____

Honor code statement: I will abide strictly by the Georgia Tech honor code at all times. I will not use a calculator. I will not reference any website, application, or other CAS-enabled service. I will not consult with my notes or anyone during this exam. I will not provide aid to anyone else during this exam.

() I attest to my integrity and will not discuss this exam with anyone until **Friday December 15**.

Read all instructions carefully before beginning.

- Print your name and GT ID neatly above.
- You may use any portion of our 2 hours and 50 minute final exam time to work on this portion of the exam.
- When you complete this portion of the exam, you may turn it in and work on another portion of the exam.
- You may not use aids of any kind.
- Please show your work.
- Good luck! Write yourself a message of encouragement on the front page!

Question	Points
1	2
2	2
3	3
4	3
5	14
6	8
7	8
Total:	40

For problems 1-3 choose whether each statement is true or false. If the statement is *always* true, pick true. If the statement is *ever* false, pick false. Be sure to neatly fill in the bubble corresponding to your answer choice.

1. (2 points) Every smooth vector field $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is conservative.

TRUE

FALSE

2. (2 points) If $\nabla \cdot \mathbf{F} = 0$ and $\nabla \times \mathbf{F} = \mathbf{0}$, then $\mathbf{F} = \mathbf{0}$.

TRUE

FALSE

3. (3 points) Let $\mathbf{F}(x, y) = \langle 2023y, -2023x \rangle$ and C be a simple closed curve surrounding the origin with positive orientation. Which of the theorems below would be appropriate to use to compute the flow of \mathbf{F} along C ?

Fundamental Theorem of Line Integrals

Green's Theorem (circulation)

Green's Theorem (flux)

Stokes' Theorem

Divergence Theorem

4. (3 points) Let $\mathbf{F}(x, y, z) = \langle 4x, -4y, 4z \rangle$ and S be the surface which is the part of the cylinder $x^2 + z^2 = 4$ between $y = 2023$ and $y = 2024$, oriented away from the y -axis. \mathbf{F} is the curl of a vector field \mathbf{G} . Which of the theorems below would be appropriate to use to compute the flux of \mathbf{F} across S ?

Fundamental Theorem of Line Integrals

Green's Theorem (circulation)

Green's Theorem (flux)

Stokes' Theorem

Divergence Theorem

5. (14 points) Let $\mathbf{F}(x, y) = \langle 2y - 3x^2, 3 \rangle$ and $\mathbf{G}(x, y) = \langle y^2 e^{xy}, (1 + xy)e^{xy} \rangle$. In this problem you will work with these vector fields and the curve C that is the portion of the parabola $y = 9 - x^2$ starting at $(-3, 0)$ and ending at $(1, 8)$.

(a) Is \mathbf{F} conservative? If so, find a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ so that $\mathbf{F} = \nabla f$.

(b) Is \mathbf{G} conservative? If so, find a function $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ so that $\mathbf{G} = \nabla g$.

(c) Compute the work done by \mathbf{F} along the curve C . Fully simplify your answer.

(d) Compute the work done by \mathbf{G} along the curve C . Fully simplify your answer.


6. Let D be the sphere of radius 3 centered at the origin in \mathbb{R}^3 . The volume of this sphere is 36π . Suppose that the density of a liquid filling this sphere is $\delta(x, y, z) = 5y^2$ kilograms per cubic meter.

(a) (4 points) Write an integral expression for the mass of this sphere filled with liquid. **Do not evaluate your integral expression.**

(b) (4 points) Write an integral expression for the average distance of a point in this sphere from the origin. **Do not evaluate your integral expression.**

7. Consider the volume D bounded by the planes $x = 0, x = 2, z = -y$ and the surface $z = y^2/2$.

- (a) (4 points) Write an integral for the volume of D using Cartesian coordinates in the order $dz dy dx$. Show your work, including a sketch of the shadow of the region. **Do not evaluate your integral.**



- (b) (4 points) Write an integral for the volume of D using Cartesian coordinates in the order $dx dy dz$. Show your work, including a sketch of the shadow of the region. **Do not evaluate your integral.**

SCRATCH PAPER - PAGE LEFT BLANK

FORMULA SHEET

- Trig identities: $\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$, $\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$
- Volume(D) = $\iiint_D dV$, $f_{avg} = \frac{\iiint_D f(x, y, z) dV}{\text{volume of } D}$, Mass: $M = \iiint_D \delta dV$
- Cylindrical coordinates: $x = r \cos(\theta)$, $y = r \sin(\theta)$, $z = z$, $dV = r dz dr d\theta$
- Spherical coordinates: $x = \rho \sin(\phi) \cos(\theta)$, $y = \rho \sin(\phi) \sin(\theta)$, $z = \rho \cos(\phi)$,
 $dV = \rho^2 \sin(\phi) d\rho d\phi d\theta$
- First moments (3D solid): $M_{yz} = \iiint_D x\delta dV$, $M_{xz} = \iiint_D y\delta dV$, $M_{xy} = \iiint_D z\delta dV$
- Center of mass (3D solid): $(\bar{x}, \bar{y}, \bar{z}) = \left(\frac{M_{yz}}{M}, \frac{M_{xz}}{M}, \frac{M_{xy}}{M}\right)$
- Substitution for double integrals: If R is the image of G under a coordinate transformation $\mathbf{T}(u, v) = \langle x(u, v), y(u, v) \rangle$ then

$$\iint_R f(x, y) dx dy = \iint_G f(\mathbf{T}(u, v)) |\det D\mathbf{T}(u, v)| du dv.$$

- Scalar line integral: $\int_C f(x, y, z) ds = \int_a^b f(\mathbf{r}(t)) |\mathbf{r}'(t)| dt$
- Tangential vector line integral: $\int_C \mathbf{F} \cdot \mathbf{T} ds = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt$
- Normal vector line integral: $\int_C \mathbf{F}(x, y) \cdot \mathbf{n} ds = \int_C P dy - Q dx = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \langle y'(t), -x'(t) \rangle dt$.
- Fundamental Theorem of Line Integrals: $\int_C \nabla f \cdot d\mathbf{r} = f(B) - f(A)$ if C is a path from A to B
- Curl (Mixed Partials) Test: $\mathbf{F} = \nabla f$ if $\text{curl } \mathbf{F} = \mathbf{0} \Leftrightarrow P_z = R_x, Q_z = R_y$, and $Q_x = P_y$.
- $\nabla = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\rangle$ $\text{div } \mathbf{F} = \nabla \cdot \mathbf{F}$ $\text{curl } \mathbf{F} = \nabla \times \mathbf{F}$
- Green's Theorem: If C is a simple closed curve with positive orientation and R is the simply-connected region it encloses, then

$$\int_C \mathbf{F} \cdot \mathbf{T} ds = \iint_R (\nabla \times \mathbf{F}) \cdot \mathbf{k} dA \qquad \int_C \mathbf{F} \cdot \mathbf{n} ds = \iint_R (\nabla \cdot \mathbf{F}) dA.$$

- Surface Area = $\iint_S 1 d\sigma$
- Scalar surface integral: $\iint_S f(x, y, z) d\sigma = \iint_R f(\mathbf{r}(u, v)) |\mathbf{r}_u \times \mathbf{r}_v| dA$
- Flux surface integral: $\iint_S \mathbf{F} \cdot \mathbf{n} d\sigma = \iint_S \mathbf{F} \cdot d\boldsymbol{\sigma} = \iint_R \mathbf{F}(\mathbf{r}(u, v)) \cdot (\mathbf{r}_u \times \mathbf{r}_v) dA$
- Stokes' Theorem: If S is a piecewise smooth oriented surface bounded by a piecewise smooth curve C and \mathbf{F} is a vector field whose components have continuous partial derivatives on an open region containing S , then

$$\int_C \mathbf{F} \cdot \mathbf{T} ds = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} d\sigma.$$

- Divergence Theorem: If S is a piecewise smooth closed oriented surface enclosing a volume D and \mathbf{F} is a vector field whose components have continuous partial derivatives on D , then

$$\iint_S \mathbf{F} \cdot \mathbf{n} d\sigma = \iiint_D \nabla \cdot \mathbf{F} dV.$$

SCRATCH PAPER - PAGE LEFT BLANK