

MATH 2551-D MIDTERM 3
VERSION A
FALL 2024
COVERS SECTIONS 15.5-8, 16.1-8

Full name: _____

GT ID: _____

Honor code statement: I will abide strictly by the Georgia Tech honor code at all times. I will not use a calculator. I will not reference any website, application, or other CAS-enabled service. I will not consult with my notes or anyone during this exam. I will not provide aid to anyone else during this exam.

() **Initial here to attest to your integrity.**

Read all instructions carefully before beginning.

- Print your name and GT ID neatly above.
- You have 75 minutes to take the exam.
- You may not use aids of any kind.
- Show your work. Answers without work shown will receive little or no credit.
- The Learning Outcomes for the exam are listed on the back of this front cover.
- The Formula Sheet is on the final page and may be removed. Do not remove any other pages. Do not write any work on the Formula Sheet
- Good luck! Write yourself a message of encouragement on the front page!

Question:	1	2	3	4	5	6	7	8	9	10	11	Total
Points:	1	1	3	3	3	3	3	3	10	10	10	50

Learning Outcomes

- **I3: Polar, Cylindrical, and Spherical Integrals.** Use polar, cylindrical, and spherical coordinates to transform double and triple integrals. Sketch regions based on given polar, cylindrical, and spherical iterated integrals.
- **I4: Triple Integrals.** Set up triple integrals as iterated integrals over any region. Sketch regions based on a given iterated integral. Interpret geometric meanings of triple integrals.
- **I5: Integral Applications.** Use multiple integrals to solve physical problems, such as finding area, average value, volume, mass of a lamina, or solid.
- **I6: Change of Variables.** Use general change of variables to transform double and triple integrals for easier calculation. Choose the most appropriate coordinate system to evaluate a specific integral.
- **V1: Scalar Line Integrals.** Define line integrals of a scalar function over a smooth curve. Evaluate scalar line integrals in two and three dimensions.
- **V2: Vector Line Integrals.** Define line integrals of a vector-valued function or vector field over a smooth curve. Evaluate vector line integrals in two and three dimensions.
- **V3: Conservative Vector Fields.** Test for conservative vector fields. Find potential functions. Evaluate line integrals using the Fundamental Theorem of Line Integrals.
- **V4: Green's Theorem.** Compute two-dimensional curls and divergence of vector fields. Evaluate integrals using Green's theorem. Solve applied problems using Green's Theorem, such as finding area.
- **V5: Surface Integrals.** Define and compute surface integrals for scalar and vector valued functions.
- **V6: Applications of Vector Calculus.** Interpret work, flow, flux, and surface area in terms of line and/or surface integrals, as appropriate.
- **V7: Generalizations of the FTC.** State and apply Stokes' Theorem and the Divergence Theorem to solve problems in three dimensions.

For problems 1-2 choose whether each statement is true or false. If the statement is *always* true, pick true. If the statement is *ever* false, pick false. Be sure to neatly fill in the bubble corresponding to your answer choice.

1. (1 point) [**V6: Applications of Vector Calculus**] If $\mathbf{F} \cdot \mathbf{T} > 0$ at every point on a curve C , then the net flow of \mathbf{F} along the curve C is positive.

TRUE

FALSE

2. (1 point) [**V3: Conservative Vector Fields**] Let \mathbf{F} be a conservative vector field in \mathbb{R}^3 . If f and g both satisfy $\nabla f = \nabla g = \mathbf{F}$, then $f = g$.

TRUE

FALSE

3. (3 points) [**V3, V4, V7**] Let $\mathbf{F}(x, y, z) = \langle 4x, -8y, 4z \rangle$ and S be the surface which is the part of the cylinder $x^2 + z^2 = 4$ between $y = 2024$ and $y = 2025$, oriented away from the y -axis. \mathbf{F} is the curl of a vector field \mathbf{G} since $\nabla \cdot \mathbf{F} = 0$. Select all of the integration theorems below that are appropriate to use to compute the flux of \mathbf{F} across S .

A) Fundamental Theorem of Line Integrals

B) Green's Theorem (circulation)

C) Green's Theorem (flux)

D) Stokes' Theorem

E) Divergence Theorem

4. (3 points) [**V3, V4, V7**] Let $\mathbf{F}(x, y) = \langle 3x + y, x - 3y \rangle$ and C be the circle $x^2 + y^2 = 25$ oriented clockwise. Select all of the integration theorems below that are appropriate to use to compute the circulation of \mathbf{F} along C .

A) Fundamental Theorem of Line Integrals

B) Green's Theorem (circulation)

C) Green's Theorem (flux)

D) Stokes' Theorem

E) Divergence Theorem

5. (3 points) [**I6: Change of Variables**] Suppose we wish to evaluate the integral $\iint_R f(x, y) dA$ using the change of variables

$$u = 3x \quad v = y^5.$$

Which of the following is the new area element dA in terms of du and dv ?

- A) $dA = 15v^{4/5} du dv$
- B) $dA = 15v^4 du dv$
- C) $dA = \frac{1}{15}v^{-4/5} du dv$
- D) $dA = du dv$
- E) $dA = \frac{5}{3}v^{-4/5} du dv$

6. (3 points) [**V3: Conservative Vector Fields**] If possible, find a potential function for the vector field

$$\mathbf{F}(x, y) = \left\langle \frac{-y}{x^2 + y^2}, \frac{x}{x^2 + y^2} \right\rangle.$$

- A) $f(x, y) = \arctan(y/x)$
- B) $f(x, y) = \arctan(y/x) + \arctan(x/y)$
- C) $f(x, y) = -\frac{1}{2} \ln(x^2 + y^2)$
- D) $f(x, y) = \frac{1}{2} \ln(x^2 + y^2)$
- E) This vector field is not conservative.

7. (3 points) [**I3: Polar, Cylindrical, and Spherical Integrals**] Which integral below computes the volume of the region inside the sphere $x^2 + y^2 + z^2 = 4$ and outside the cylinder $x^2 + y^2 = 2$?

- A)** $\int_0^{2\pi} \int_{\sqrt{2}}^2 \int_0^{\sqrt{4-r^2}} dz dr d\theta$
 B) $\int_0^\pi \int_0^2 \int_{-\sqrt{4-r^2}}^{\sqrt{4-r^2}} r dz dr d\theta$
 C) $\int_0^{2\pi} \int_0^{\pi/4} \int_{\sqrt{2} \csc(\phi)}^2 \rho^2 \sin(\phi) d\rho d\phi d\theta$
 D) $\int_0^{2\pi} \int_{\pi/4}^{3\pi/4} \int_0^2 \rho^2 \sin(\phi) d\rho d\phi d\theta$
 E) $\int_0^{2\pi} \int_{\pi/4}^{3\pi/4} \int_{\sqrt{2} \csc(\phi)}^2 \rho^2 \sin(\phi) d\rho d\phi d\theta$

8. (3 points) [**V5: Surface Integrals**] Select all of the parameterizations $\mathbf{r}(s, t)$ below corresponding to the surface which is the part of the elliptical paraboloid $x = y^2 + 4z^2$ with $0 \leq x \leq 4$.

- A)** $\mathbf{r}(s, t) = \langle s, t, s^2 + 4t^2 \rangle, \quad 0 \leq s^2 + 4t^2 \leq 4$
 B) $\mathbf{r}(s, t) = \langle s^2 + 4t^2, s, t \rangle, \quad 0 \leq s^2 + 4t^2 \leq 4$
 C) $\mathbf{r}(s, t) = \langle s \cos(t), s \sin(t), s^2 \cos^2(t) + 4s^2 \sin^2(t) \rangle, \quad 0 \leq s \leq 4, 0 \leq t \leq 2\pi$
 D) $\mathbf{r}(s, t) = \langle s, \sqrt{s} \cos(t), \frac{\sqrt{s}}{2} \sin(t) \rangle, \quad 0 \leq s \leq 4, 0 \leq t \leq 2\pi$
 E) $\mathbf{r}(s, t) = \langle 4t^2, 2t \cos(s), t \sin(s) \rangle, \quad 0 \leq s \leq 2\pi, 0 \leq t \leq 1$

9. (10 points) [**I4: Triple Integrals, I5: Integral Applications**] Set up but do not compute an integral to compute the mass of the solid D that has mass density function $\delta(x, y, z) = 4 + \cos(x) \sin(y) + z/10$ and occupies the region

$$y \leq 8 - x^2, \quad x^2 + 2z^2 \leq y, \quad x \geq 0.$$

A complete solution will include a sketch of the shadow of D in the plane of the two variables that you do not integrate first.

10. (10 points) [**V5: Surface Integrals, V6: Applications of Vector Calculus, V7: Generalizations of the FTC.**] Compute the flux of the vector field

$$\mathbf{F}(x, y, z) = \left\langle xy^2 + \frac{1}{3}x^3, yx^2 + e^{x \cos(z)}, y^2z + \sin(\arctan(x)) \right\rangle$$

out of the closed surface S which is the portion of the paraboloid $z = x^2 + y^2$ from $z = 0$ to $z = 4$ together with the disk $x^2 + y^2 \leq 4$ in the plane $z = 4$. Simplify your final answer.

Hint: Cylindrical coordinates may be useful.

11. [**V2: Vector Line Integrals, V3: Conservative Vector Fields, V4: Green's Theorem**] Consider the vector field $\mathbf{F} = \langle 2y, x \rangle$ and the curve C which is the circle $x^2 + y^2 = 25$, oriented counterclockwise with outward normal vector.

(a) (4 points) Compute $\int_C \mathbf{F} \cdot \mathbf{T} \, ds$.

(b) (4 points) Compute $\int_C \mathbf{F} \cdot \mathbf{n} \, ds$.

(c) (2 points) Based on your work above, what can you conclude about whether \mathbf{F} is conservative? Justify your answer.

SCRATCH PAPER - PAGE LEFT BLANK

SCRATCH PAPER - PAGE LEFT BLANK

FORMULA SHEET

Rules for Triple Integrals for the Sketching Impaired

- 1: Choose a variable appearing exactly twice for the next integral.
- 2: After setting up an integral, cross out any constraints involving the variable just used.
- 3: Create a new constraint by setting the lower limit of the preceding integral less than the upper limit.
- 4: A squared variable counts twice.
- 5: The argument of a square root must be nonnegative.

- Trig identities: $\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$, $\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$
- Volume(D) = $\iiint_D dV$, $f_{avg} = \frac{\iiint_D f(x, y, z) dV}{\text{volume of } D}$, Mass: $M = \iiint_D \delta dV$
- Coordinate Transforms:
 - Cylindrical: $x = r \cos(\theta)$, $y = r \sin(\theta)$, $z = z$, $dV = r dz dr d\theta$
 - Spherical: $x = \rho \sin(\phi) \cos(\theta)$, $y = \rho \sin(\phi) \sin(\theta)$, $z = \rho \cos(\phi)$,
 $dV = \rho^2 \sin(\phi) d\rho d\phi d\theta$
- Substitution Theorem: If R is the image of G under a coordinate transformation $\mathbf{T}(u, v) = \langle x(u, v), y(u, v) \rangle$ then

$$\iint_R f(x, y) dx dy = \iint_G f(\mathbf{T}(u, v)) |\det D\mathbf{T}(u, v)| du dv.$$

- Line integrals:
 - $\int_C f(x, y, z) ds = \int_a^b f(\mathbf{r}(t)) \|\mathbf{r}'(t)\| dt$
 - $\int_C \mathbf{F} \cdot \mathbf{T} ds = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt$
 - $\int_C \mathbf{F} \cdot \mathbf{n} ds = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \langle y'(t), -x'(t) \rangle dt.$
- Fundamental Theorem of Line Integrals: $\int_C \nabla f \cdot d\mathbf{r} = f(B) - f(A)$ if C is a path from A to B
- Curl (Mixed Partial) Test: On a simply-connected region $\mathbf{F} = \nabla f \Leftrightarrow \text{curl } \mathbf{F} = \mathbf{0}$
- $\nabla = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right\rangle$ $\text{div } \mathbf{F} = \nabla \cdot \mathbf{F}$ $\text{curl } \mathbf{F} = \nabla \times \mathbf{F}$
- Green's Theorem: If C is a simple closed curve with positive orientation and R is the simply-connected region it encloses, then

$$\int_C \mathbf{F} \cdot \mathbf{T} ds = \iint_R (\nabla \times \mathbf{F}) \cdot \mathbf{k} dA \qquad \int_C \mathbf{F} \cdot \mathbf{n} ds = \iint_R (\nabla \cdot \mathbf{F}) dA.$$

FORMULA SHEET CONTINUED

- Surface integrals:

$$- \iint_S f(x, y, z) \, d\sigma = \iint_R f(\mathbf{r}(u, v)) \|\mathbf{r}_u \times \mathbf{r}_v\| \, dA$$

$$- \iint_S \mathbf{F} \cdot \mathbf{n} \, d\sigma = \iint_S \mathbf{F} \cdot d\boldsymbol{\sigma} = \iint_R \mathbf{F}(\mathbf{r}(u, v)) \cdot (\mathbf{r}_u \times \mathbf{r}_v) \, dA$$

- Stokes' Theorem: If S is a piecewise smooth oriented surface bounded by a piecewise smooth curve C and \mathbf{F} is a vector field whose components have continuous partial derivatives on an open region containing S , then

$$\int_C \mathbf{F} \cdot \mathbf{T} \, ds = \iint_S (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\sigma.$$

- Divergence Theorem: If S is a piecewise smooth closed oriented surface enclosing a volume D and \mathbf{F} is a vector field whose components have continuous partial derivatives on D , then

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, d\sigma = \iiint_D \nabla \cdot \mathbf{F} \, dV.$$